

Lacitone

Furosemide & Spironolactone

Presentation:-

Each film coated tablet contains Furosemide BP 20 mg and Spironolactone BP 50 mg.

Indications:-

Lacitone is indicated for -

- Essential hypertension
- Congestive heart failure
- Liver cirrhosis with ascites
- Swelling due to excess fluid retention (edema)
- Hyperaldosteronism

Dosage and Administration:-

1 to 4 tablets daily, (20 to 80 mg of Furosemide and 50 to 200 mg of Spironolactone) according to patient's response.

Contraindications:-

Lacitone is contraindicated in patient with hyperkalemia, hyponatremia, acute renal insufficiency and hypersensitivity to Furosemide or to Spironolactone or to any ingredient containing this product.

Side Effects:-

Fatigue, nausea, malaise, gastric upset, headache, hypotension, skin rash.

Precautions:-

Electrolyte imbalance should be monitored. Special precaution should be taken in patients of diabetes, prostatic hypertrophy, hypotension, hypovolemia.

Use in pregnancy and lactation:-

Lacitone should be used with caution during pregnancy and only if expected benefit to mother is greater than the possible risk to the fetus. Furosemide passes into breast milk. It's not recommended while breast feeding.

Drug interactions:-

Drug interaction may occur with the following drugs: ACE inhibitors, potassium salts, Cardiac glycosides, Corticosteroids, NSAIDs.

Commercial pack:-

Lacitone: Each box contains 3 blister packs of 10 tablets.

Manufactured by :

 **GENERAL**
Pharmaceuticals Ltd.
Gazipur, Bangladesh



Revision No.: 00