

Tamoxen

(Tamoxifen B.P)

Presentation :

Tamoxen tablet 10mg : Each film-coated tablet contains Tamoxifen BP 10mg.

Tamoxen tablet 20mg : Each film-coated tablet contains Tamoxifen BP 20mg.

Indications :

Tamoxifen is used in treatment of

1. Breast cancer
 - a. As an adjuvant immediately after surgery in patient with early disease
 - b. For advanced breast cancer
2. Infertility.

Dosage and administration :

In breast cancer 20mg daily; anovulatory infertility, 20mg daily on days 2, 3, 4 and 5 of cycle; if necessary the dose may be increased to 40mg then 80mg for subsequent courses; if cycles irregular, start initial course on any day, with subsequent course starting 45 days later or on day 2 of cycle if menstruation occurs.

Precautions :

Occasional cystic ovarian swellings in premenopausal women, occasional hypercalcemia if bony metastases; breast-feeding; endometrial changes; porphyria; increased risk of thromboembolic events when used with cytotoxics.

Side-effects :

Hot flushes, vaginal bleeding and vaginal discharge, suppression of menstruation in some premenopausal women, pruritus vulvae, gastro-intestinal disturbances, headache, light-headedness, tumour flare, decreased platelet counts; occasionally oedema, alopecia, rashes, uterine fibroids; also visual disturbances; leucopenia, rarely neutropenia; hypertriglyceridaemia; thromboembolic events; liver enzyme changes; rarely interstitial pneumonia, hypersensitive reactions including angioedema, Steven-Johnson syndrome, bullous pemphigoid etc are reported.

Contraindications :

Tamoxifen is contraindicated in Pregnancy.

Drug interactions :

Tamoxifen has been reported to considerably increased the effect of warfarin, with possibly fatal result. It causes hepatotoxicities when given concurrently with allopurinol.

In pregnancy and lactation :

Tamoxifen should not be used in pregnancy and lactation.

Commercial pack :

Tamoxen tablet 10mg : Each box contains 3x10' tablets in blister pack.

Tamoxen tablet 20mg : Each box contains 3x10' tablets in blister pack.



GENERAL Pharmaceuticals Ltd.

Gazipur, Bangladesh.