

Genamox

Amoxicillin Capsule/Suspension

Presentation :

Genamox 500 mg Capsule : Each capsule contains Amoxicillin Trihydrate BP equivalent to 500 mg Amoxicillin.

Genamox 250 mg Capsule : Each capsule contains Amoxicillin Trihydrate BP equivalent to 250 mg Amoxicillin.

Genamox powder for suspension : After reconstitution each 5ml suspension contains Amoxicillin Trihydrate BP equivalent to 125 mg Amoxicillin.

Genamox powder for Paediatric Drops : After reconstitution each 1.25 ml suspension contains Amoxicillin Trihydrate BP equivalent to 125 mg Amoxicillin.

Indications :

Genamox (Amoxicillin) is a broad-spectrum antibiotic. Amoxicillin is indicated for the treatment of the following bacterial infections that is occurred by both gram-positive and gram-negative organism.

Respiratory tract and ENT infections : Acute and chronic bronchitis, pneumonia, otitismedia, sinusitis, tonsillitis, pharyngitis and laryngitis.

Urinary tract infections : Pyelonephritis, cystitis and urethritis.

Obstetric and gynecological infections : Bacteriuria in pregnancy, septic abortion, abdominal sepsis and puerperal sepsis.

Gastro-intestinal infections : Typhoid and paratyphoid.

Skin and soft tissue infections : Cellulitis, infected wounds and abscesses.

Generalized infections : Septicemia, bacterial endocarditis, meningitis, peritonitis and osteomyelitis.

Sexual transmitted diseases : Gonorrhoea and syphilis.

Amoxicillin may also be used as prophylactic cover for patients at risk of developing endocarditis when undergoing surgery.

Dose and Administration :

Adult : 250-500 mg 8 hourly 3 times daily.

Children : 2 to 10 years, 125 mg every 8 hours. Double in severe infections.

Children below 2 years : $\frac{1}{4}$ of adult dose.

Contraindications :

Amoxicillin is contraindicated to the patients of Penicillin hypersensitivity, Glandular fever and lymphatic lymphoma, Bacterial resistance.

Side Effects :

Side effects are usually of mild transitory nature. They are most likely to occur diarrhea, dyspepsia, allergy, amoxicillin should be discontinued when any types of skin rash occur.

Special precautions for storage :

The reconstituted suspension must be used within 7 days if kept at room temperature and within 10 days when stored in a refrigerator.

Commercial Pack :

Genamox 500 mg Capsule : Each box containing 5 blister packs of 10 capsules.

Genamox 250 mg Capsule : Each box containing 10 blister packs of 10 capsules.

Genamox powder for suspension : Each bottle contains dry powder for reconstitution into 100 ml suspension.

Genamox powder for Paediatric Drops : Each bottle contains dry powder for reconstitution into 15 ml suspension.

 **GENERAL**
Pharmaceuticals Ltd.
Gazipur, Bangladesh