

# DOMILIN

Domperidone maleate BP

**Presentation :**

**Domilin Tablet :** Each tablet contains Domperidone maleate BP equivalent to 10 mg Domperidone.

**Domilin suspension :** Each 5 ml contains 5 mg Domperidone.

**Description :** Domperidone is a benzimidazole derivative which has prokinetic and antiemetic properties.

**Indication & uses :**

Symptomatic treatment of the dyspeptic symptom complex which is often associated with delayed gastric emptying or gastroesophageal reflux and

**oesophagitis :** epigastric sense of fullness, feeling of abdominal distension, upper abdominal pain, flatulence, heartburn. Treatment of nausea and vomiting of various origins including functional organic, infectious, dietic, radiotherapy, medicine therapy.

Domilin relieves the nausea associated with migraine attacks.

**Dosage & Administration :**

Chronic dyspepsia :

**Adults :** 10 mg (1 Tab ) 3 times daily, 15-30 minutes before meals, if necessary it can be repeated before retiring.

**Children :** 0.2-0.4 mg/kg every 4-8 hours daily before meals, if necessary, it can be repeated before retiring.

Na usea and Vomiting :

**Adults :** 10-20 mg, 3-4 times daily before meals and before retiring.

**Children :** 0.2-0.4 mg/kg every 4-8 hours daily before meals and before retiring.

**Contraindication :**

❖ Patients with intolerance to this drug.

❖ Do not use it the increase in gastric motility can be dangerous eg. bleeding, mechanical obstruction or gastrointestinal perforation.

**Side effects :**

The most frequent reactions to Domperidone are those related to elevated prolactin levels including breast tenderness, galactorrhoea, gynaecomastia and amenorrhoea. This effects are dose-related and gradually resolve after lowering the dose or discontinuing treatment. Other rarely reported adverse reactions include headache, diarrhoea, dizziness, mild and transient abdominal cramps, dry mouth and drowsiness. Rare allergic reactions such as rash and urticaria have also been reported.

**Precaution & Warning :**

❖ Do not give antacids or antisecretory drugs concomitantly with domperidone. If the administration of these drugs is required. Domilin must be given after meals. Exercise caution in patient with liver function disturbances.

❖ Use in children : Blood brain barrier and metabolism functions in babies have not been fully developed yet, therefore the possibility of neurologic side effect can not be avoided in infants <1 year.

**Interactions :**

Anticholinergic drugs may inhibit antidyspeptic effect of domperidone. Antacid and antisecretory drugs can reduce bioavailability of domperidone if administered together.

**Commercial pack :**

**Domilin Tablet :** Box contains 10 blister strips of 10 tablets.

**Domilin Suspension :** Each bottle contains 100 ml suspension



**GENERAL Pharmaceuticals Ltd.**

Gazipur, Bangladesh.