

Travagen Z

Travoprost
Sterile Eye Drops

Presentation

Travagen Z: Each ml sterile eye drops contains Travoprost USP 0.04 mg.

Preservative: Ionic Buffered System

Pharmacological Action

Travoprost free acid is a selective FP prostanoid receptor agonist which is believed to reduce intraocular pressure by increasing uveoscleral outflow. The exact mechanism of action is unknown at this time.

Indications

Travoprost decreases elevated intraocular pressure (IOP) in adult patients with open-angle glaucoma or ocular hypertension who are intolerant of or insufficiently responsive to another intraocular pressure lowering medication as monotherapy or as adjunctive therapy.

Dosage and Application

The recommended dosage is one drop of Travoprost in the conjunctival sac of the affected eye(s) once daily in the evening. If more than one topical ophthalmic product is being used, the mentioned medicine should be administered at least five minutes apart. While substituting another ophthalmic anti-glaucoma agent with Travoprost, discontinue the other agent and start the following day with Travoprost.

Contraindications

Contraindicated in patients with hypersensitivity to any component of this preparation.

Side Effects

Ocular hyperemia is the most frequently reported treatment related side effect.

Precautions

Travoprost should be used with cautions in patients with active intraocular inflammation (iritis/uveitis). Contact lenses should be removed prior to the administration of the medicine. Lenses may be reinserted 15 minutes after administration of Travoprost.

Warning

Travoprost may cause changes to pigmented tissues.

Use in specific population

Pregnancy

There are no adequate data from the use of Travoprost in pregnant women. The potential risk for humans is unknown.

Lactation

It is not known whether Travoprost is excreted in human breast milk. Cautions should be exercised during administration of Travoprost to a lactating mother.

Drug Interactions

No interaction studies have been performed with Travoprost.

Over Dose

No data are available with regard to overdose with this eye drops.

Storage Conditions

Store between 2°-25°C and away from light. Keep out of reach of children. Do not touch the dropper tip to surfaces since this may contaminate the solution. After one month of the first opening do not use the medicine of dropper.

Commercial Pack

Travagen Z: Each plastic dropper bottle containing 3 ml sterile eye drops.

Manufactured by :

 **GENERAL**
Pharmaceuticals Ltd.
Unit-2, Gazipur, Bangladesh

Revision No. : 00

22060251