

Broxine Syrup

Bromhexine HCl BP

Presentation :

Broxine Syrup : Each 5 ml syrup contains Bromhexine HCl BP 4 mg.

Indications :

Broxine (Bromhexine) is used in the treatment of respiratory disorders associated with productive cough.

Dosage and Administration :

Adults : Recommended daily dose is 2 to 4 teaspoonfuls 3 times. Initially 4 teaspoonfuls 3 times daily and then as required.

Children : Suggested dosage for children under 2 years is ¼ teaspoonful 3 times daily, for 2-5 years ½ teaspoonful 3 times daily and for children aged 5-10 years 1 teaspoonful 3 times daily.

Precautions :

Since mucolytics may disrupt the gastric mucosal barrier, so Bromhexine should be used with care in patients with a history of peptic ulceration. Care is also advisable in asthmatic patients.

Contraindications :

Contraindicated to those who are hypersensitive to Bromhexine.

Side effects :

Gastro-intestinal side effects may occur occasionally with Bromhexine and a transient rise in serum aminotransferase values has been reported. Other reported side effects include headache, dizziness, sweating and skin rash.

Storage Conditions :

Store below 25°C, protect from light. Keep the container tightly closed.

Commercial pack :

Each bottle contains 100 ml syrup.

 **GENERAL**
Pharmaceuticals Ltd.
Gazipur, Bangladesh