

Infud Tablet

(Terbinafine HCl)

Presentation

Infud Tablet: Each Tablet contains terbinafine hydrochloride equivalent to Terbinafine 250 mg.

Indications

- 1) Dermatophyte infections of the nails (Onychomycosis)
- 2) Ringworm infections including tinea pedis, tinea cruris and tinea corporis where oral therapy is appropriate (due to site, severity or extent)

Dosage and Administration

250 mg daily usually for 2-6 weeks in tinea pedis, 2-4 weeks in tinea cruris, 4 weeks in tinea corporis, 6 weeks-3 months in nail infections (occasionally longer in toenail infections); Child: usually for 2 weeks, tinea capitis, over 1 year, body-weight 10-20kg, 62.5 mg once daily; body-weight 20-40 kg, 125 mg once daily; body-weight over 40 kg, 250 mg once daily.

Side-effects

Abdominal discomfort, anorexia, nausea, diarrhoea, headache, rash and urticaria occasionally with arthralgia or myalgia, serious skin reactions including Stevens-Johnson syndrome and toxic epidermal necrolysis may be reported (discontinue treatment if progressive skin rash occurs), taste disturbance, photosensitivity, and rarely liver toxicity (discontinue treatment) including , jaundice, cholestasis and hepatitis may also be reported.

Contraindications

Hypersensitivity to terbinafine or any of the excipient contained in the tablet.

Drug Interactions

Antibacterial agents: plasma concentration is increased by Rifampicin, Ulcer healing drugs : plasma concentration is reduced by Cimetidine. Occasional report of bleeding with oral contraceptives.

Use in Pregnancy and Lactation

Reproduction studies in animal do not suggest that there is any risk of fetal damage but this has not been confirmed by controlled trials in pregnant woman. Terbinafine should therefore not be given to pregnant woman unless the potential benefits clearly outweigh the risks. Terbinafine is excreted in breast milk and although the amount excreted is small, mothers taking Terbinafine should not feed breast milk.

Commercial Pack

Infud Tablet: Each box contains 1 blister strip of 4 tablets.

Manufactured by :
 **GENERAL**
Pharmaceuticals Ltd.
Gazipur, Bangladesh

