

# Deleta

Flupentixol & Melitracen Tablet

## Presentation

**Deleta:** Each film coated tablet contains Flupentixol 0.5 mg and Melitracen 10 mg (as Flupentixol Hydrochloride BP & Melitracen Hydrochloride INN).

## Indications

1. In the treatment of anxiety, depression and apathy.
2. Maintenance in psychogenic depressions and other psychoses.

## Dosage and administration

*Adult :* Usually 2 tablets daily divided in morning and evening doses.

*In severe case :* Morning dose may be increased to 2 tablets.

*Elderly :* 1 tablet in the morning.

*Children :* Not recommended.

## Precautions

If previously the patient has been treated with tranquilizers with sedative effect these should be withdrawn gradually.

## Side-effects

Restlessness, insomnia, hypomania reported, rarely dizziness, tremor, visual disturbances, headache, hyperprolactinaemia, extrapyramidal symptoms.

## Contra-indications

Extreme agitation, cardiac disorders, CV insufficiency. Patients receiving MAOIs within 14 days.

## Drug-interactions

Flupentixol and melitracen may enhance the response to alcohol, barbiturates and other CNS depressants. Simultaneous administration of MAOIs may cause hypertensive crisis. Neuroleptics and thymoleptics reduce the antihypertensive effect of guanethidine and similar acting compounds and thymoleptics enhance the effects of adrenaline and noradrenaline.

## Use in pregnancy and Lactation

Patients should avoid this drug during pregnancy and lactation.

## Commercial pack

**Deleta:** Each box contains 5 blister packs of 10 tablets.

Manufactured by :

 **GENERAL**  
Pharmaceuticals Ltd.  
Gazipur, Bangladesh

