

# LORIN

Loratadine USP

**Presentation :**

**Lorin tablet :** Each tablet contains Loratadine USP 10 mg.

**Lorin suspension :** Each 5 ml suspension contains Loratadine USP 5 mg.

**Indications :**

- ❖ Allergic rhinitis such as sneezing, nasal discharge, itching, ocular itching & burning.
- ❖ Seasonal allergic rhinitis
- ❖ Perennial allergic rhinitis
- ❖ Symptoms of hay fever
- ❖ Symptoms associated with idiopathic chronic urticaria

**Dosage & administration :**

*Adult & Child over 6 years :* 10 mg once daily.

*Child 2-5 years :* 5 mg once daily.

*Child below 2 years :* Safety and efficacy has not been established.

**Side effects :**

Fatigue, nausea and headache were rarely reported. Incidence of sedation and antimuscarinic effects were low.

**Contraindications :**

Lorin (Loratadine) is contraindicated in patients who have severe hypersensitivity or idiosyncrasy to its components.

**Interactions :**

When administered concurrently with alcohol, Loratadine has no potentiating effect as measured by psychomotor performance studies. Interaction with other drugs has not been reported.

**Warnings :**

Antihistamines should be used with caution in hepatic disease and dose reduction may be necessary in renal impairment. Caution may be required in epilepsy. It should be used with caution in urinary retention, glaucoma and pyloroduodenal obstruction.

**Pregnancy and lactation :**

Lorin (Loratadine) should not be used during pregnancy and Lactation.

**Storage :**

Lorin (Loratadine) should be stored in a cool and dry place.

**Commercial pack :**

**Lorin tablet :** Each box contains 10 blister strips of 10 tablets.

**Lorin Suspension :** Each bottle contains 60 ml Lorin suspension.