

Ologen MAX

Olopatadine 0.7%

Sterile Eye Drops

Presentation

Ologen MAX: Each ml sterile eye drops contains Olopatadine 7 mg (As Olopatadine Hydrochloride USP).

Preservative: Sodium Perborate 0.01%

Pharmacological Action

Olopatadine inhibits the release of histamine from the mast cell and is relatively selective histamine H1 receptor antagonist. Which inhibits type 1 immediate hypersensitivity reaction including inhibition of histamine induced effects on human conjunctival epithelial cells. Olopatadine has no effects on alpha-adrenergic, dopamine and muscarinic type 1 and 2 receptors.

Indications

Olopatadine eye drops are indicated for the treatment of signs and symptoms (itchy, watery, red and swollen eyes or eyelids) of allergic conjunctivitis including vernal keratoconjunctivitis, blepharitis, blepharoconjunctivitis and giant papillary conjunctivitis.

Dosage and Application

The recommended dose of Olopatadine eye drops is 1 drop in the affected eye(s) once daily.

Contraindications

It is contraindicated in persons with a known hypersensitivity to any component of this product.

Side Effects

Rarely headache have been reported. Other adverse effects: asthenia, blurred vision, burning or stinging, cold syndrome, dry eye, foreign body sensation, hyperemia, hypersensitivity, keratitis, lid edema, nausea, pharyngitis, pruritus, rhinitis, sinusitis, and taste perversion.

Precautions

Patients should be advised not to wear a contact lens if their eye is red. Olopatadine ophthalmic solution should not be used to treat contact lens related irritation. Sodium Perborate, preservative of this eye solution, may be absorbed by soft contact lenses. Patients who wear soft contact lenses and whose eyes are not red should be instructed to wait at least 10 minutes after instillation of Olopatadine eye drops before they insert their contact lenses. The treatment should be discontinued in the presence of an allergic reaction.

Drug Interactions

Specific drug interactions studies have not been conducted with Olopatadine ophthalmic solution.

Use in Pregnancy and Lactation

There are no adequate and well controlled studies in pregnant women. This drug should be used in pregnant women only if the potential benefit justifies the potential risk to the fetus.

Overdose

There is no information on Olopatadine overdose. However, excessive use of any medication can have serious consequences. If you suspect an overdose, seek medical attention without delay.

Storage Conditions

Store in a cool and dry place, away from light. Keep out of reach of children. Do not touch the dropper tip to surfaces since this may contaminate the solution. After one month of the opening do not use the medicine of dropper.

Commercial Pack

Ologen MAX: Each plastic dropper bottle containing 5 ml sterile eye drops.

Manufactured by :

 **GENERAL**
Pharmaceuticals Ltd.
Unit-2, Kaliakair, Gazipur, Bangladesh

22060125

Revision No.: 00