# Eclo

Clobetasol Propionate USP 0.05% w/w (Only for external use)

## Presentation:

**Eclo Cream :** Each 100 mg cream contains 0.05 mg of Clobetasol Propionate USP.

**Eclo Ointment**: Each 100 mg ointment contains 0.05 mg of Clobetasol Propionate USP.

#### Indication and uses:

Moderate to severe corticosteroid-responsive dermatoses. Treatment of more resistant dermatoses eg. stubborn eczemas, psoriasis, recalcitrant eczemas, lichen planus, discoid lupus, erythematosus and other conditions which do not respond satisfactorily to less active steroids.

## Dosage and Administration:

Apply sparingly 1-2 times daily for up to 4 weeks, reducing frequency as condition responds.

## Contra-indication:

Untreated bacterial, fungal, or viral skin lesions.

#### **Uses in Pregnancy and Lactation:**

Eclo is safe to use in pregnancy and lactation.

#### Side-effects:

Side effects from clobetasol use are rare; but local side effects from the use of clobetasol include: Atrophic changes, such as striae, thining of the skin; dilatation of the superficial blood vessels; spread and worsening of untreated infection; acne at the site of application in some patients.

# Warning and Precautions:

Children, especially babies, are particularly susceptible to side-effects of clobetasol. So avoid prolong clobetasol use in infants and children and on the face.

# Commercial pack:

**Eclo Cream**: 10 gm cream in aluminium collapsible tube. **Eclo Ointment**: 10 gm ointment in aluminium collapsible tube.

