

Tolcalm

(Tolperisone HCl Tablet)

Presentation :

Tolcalm : Each film-coated tablet contains Tolperisone Hydrochloride INN 50 mg.

Indications :

Treatment of increased tone of skeletal muscles due to organic neurological disorders such as injury of the pyramidal tract, multiple sclerosis, myelopathy etc.

Treatment of muscular hypertension, muscular spasm, muscular contracture, rigidity, spinal automatism and discopathy.

In the treatment of obliterative vascular disease such as obliterative arteriosclerosis, diabetic angiopathy, obliterative thromboangitis, Raynaud's disease, diffuse scleroderma etc.

Treatment of disorders due to injured innervation of the vessels such as acrocyanosis, intermittent angioneurotic dysbasia etc.

Dosage and administration :

The recommended dose is as follows;

Adult : 50-150 mg three times daily according to the individual requirement and tolerance of the patient.

Children (from 3 months to 6 years) : 5-10 mg/kg/day in three divided doses.

Children (from 6 to 14 years) : 2-4 mg/kg/day in three divided doses.

Contraindications :

Hypersensitivity to any component of the product, Myasthenia gravis, Nursing mother.

Precautions :

In case of children, the prescribed dose and duration of treatment should closely be observed.

Side effects :

Sometimes muscular weakness, headache, nausea, vomiting and rarely hypersensitivity reactions may occur as side effects.

Use in pregnancy and Lactation :

Tolperisone hydrochloride should be used in pregnancy according to physician's advice. Tolperisone hydrochloride should not be used during breast feeding.

Drug interactions :

Concomitant use of methocarbamol with tolperisone hydrochloride has been reported to cause disturbance of visual accommodation.

Packaging Quantities :

Each box contains 5X10's /10X10's tablets in blister pack.



GENERAL Pharmaceuticals Ltd.

Gazipur, Bangladesh.