

# TRICODERMA

Econazole Nitrate + Triamcinolone Acetonide

(Only for external use)

**Description :** Tricoderma is a strong antifungal, antibacterial and steroid combined preparations. It is white in color and contains Econazole Nitrate BP 1% and Triamcinolone Acetonide BP 0.1%.

**Indications :** Triamcinolone Acetonide is a strong effective topical steroid and has quick effect as an anti inflammatory, anti pruritic and anti allergic action.

Econazole Nitrate is a broad spectrum antifungal drug. It is potent, active against various dermatophyte, yeast and mould. Besides these it has antibacterial activity against gram positive microorganism.

- Tricoderma is indicated in the dermatomycoses caused by the dermatophyte, yeast and fungus.
- It is used for the management of signs and symptoms of certain inflammation and allergic conditions such as eczematous mycoses, diapa dermatitis, eczema marginatum, intertrigo, folliculitis, trichophytica and sycosis barbae.
- It is used in the treatment of different types of mycoses in various fold of body in where intolerate any drug.

**Dosage & Application :** The affected areas are washed and dry thoroughly. Then sufficient amount of cream gently massages with the finger, once or twice in a day in the morning or in the evening. This cream should be applied for at least two weeks after complete disappear of all signs of infections. Do not occlusive dressing with tricoderma during the treatment. The treatment should be stopped after 3-4 weeks use with this drug as it impaired the skin like other steroid drug.

**Side effects :** Skin irritation, local burning rarely and skin rash.

**Contraindications :** It is contraindicated like other steroid drug in TB, luetic and in viral skin infections such as herpes, vaccinia, vericella. In pregnancy no evidence has been established in human being's. It should not be used on pregnant woman during first trimester for long time or overdose.

**Warnings :** The prolong use of Tricoderma should be avoided in the thin skin and face. Duration of treatment is 2-4 weeks enough for complete recovery for eczematous but it should be continued for the disappearance of inflammatory mycoses. It will have to care in the eczema chroris vericosam so that Tricoderma does not touch in ulcus proper.

**Precautions :** It should be avoided for prolong use in topical corticosteroid fo r children, as there is a risk for adrenal suppression. Hypersensitivity is occurred in rare case if it is happened the therapy should be stopped.

**Commercial pack :** 10gm cream in aluminium collapsible tube.



**GENERAL Pharmaceuticals Ltd.**

Gazipur, Bangladesh.